



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

LONG-COVID: COGNITIVE DEFICITS (BRAIN FOG) AND BRAIN LESIONS IN NON-HOSPITALIZED PATIENTS

Jacques HUGON

PII: S0755-4982(21)00023-3
DOI: <https://doi.org/10.1016/j.lpm.2021.104090>
Reference: LPM 104090



To appear in: *La Presse Médicale*

Please cite this article as: Jacques HUGON , LONG-COVID: COGNITIVE DEFICITS (BRAIN FOG) AND BRAIN LESIONS IN NON-HOSPITALIZED PATIENTS, *La Presse Médicale* (2021), doi: <https://doi.org/10.1016/j.lpm.2021.104090>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Elsevier Masson SAS. All rights reserved.

LONG-COVID: COGNITIVE DEFICITS (BRAIN FOG) AND BRAIN LESIONS IN NON-HOSPITALIZED PATIENTS

Jacques HUGON

Centre de Neurologie Cognitive Université de Paris AHP Hôpital Lariboisière
75010 Paris France

Clinique de la Mémoire 16 rue de Téhéran 75008 Paris France

jacques.hugon@inserm.fr

The COVID-related pandemic has affected tens of millions of people around the world and resulted in a large number of hospitalizations in either intensive care or in the conventional sector (1). The initial target of Sars-Cov-2 is the upper and lower respiratory tract leading to respiratory failure requiring oxygen therapy or mechanical ventilation. Other organs can be involved in COVID such as the heart, the digestive system and blood vessels during direct viral diffusion or through a delayed inflammatory and cytokine storm (2). Clinical forms can be life-threatening, while in other cases multiple organ lesions are observed following the acute phase in intensive care units or after regular hospitalization. The brain can also be directly or indirectly affected during Sars-Cov-2 infection. A few cases of direct viral cerebral infection have been reported in acute encephalitis. The virus has rarely been detected in the cerebrospinal fluid, a finding suggesting a possible invasion via the olfactory tract (anosmia is common) or by rupture of the blood brain barrier (2). The few published neuropathological studies of encephalitis cases mainly found ischemic vascular damage, hemorrhagic or inflammatory

lesions, sometimes with molecular evidence of viral infections (3, 4). Cerebrospinal fluid anomalies confirming neuronal and astrocytic demise have also been noted, including increased concentrations of neurofilaments and GFAP (5).

In a recent study of 236,379 patients with COVID, the incidence of neurological and psychiatric disorders at 6 months was 33.6% (6). Many neuropsychiatric conditions have been described after infection with SARS-CoV-2 including chronic malaise, diffuse myalgia, headache, anosmia, cognitive impairment, depression and anxiety (7-9).

If neurological complications are more frequent in hospitalized patients in intensive care units or in regular wards, neurological disorders, particularly cognitive disorders, have also been reported in patients who had mild to moderate forms of COVID infections without hospital admission (10). A study from the Chicago Neuro-COVID Center included 50 COVID-positive patients and 50 COVID-negative (not tested) patients over a 6-month period and compared long-term neurological and cognitive consequences in both groups. All 100 patients had signs of COVID infection and were seen on average 4.7 months after the onset of infection. No patient was hospitalized. The mean age of the patients was 43.2 years. Signs of depression and anxiety were found in 42% of infected cases. 72% of the patients were females. The most common neurological symptoms were cognitive impairment with brain fog (81%), headaches (68%), paresthesia (60%), ageusia (59%), anosmia (55%), myalgia (55%) dizziness (47%) and pain (43%). Chronic fatigue and impaired daily living activities were reported in 85% of patients. Brain MRIs were found abnormal in 18% of tested patients, mainly revealing white matter changes. COVID-positive patients had more impaired

performance on attentional and working memory tests. The main conclusion of this study was that following mild COVID infections, cognitive symptoms were more frequent in young females than in young males.

The brain FDG PET scan can be very useful as a means detecting post-COVID brain damage. A recent study showed that in two hospitalized patients who had a COVID infection, brain FDG PET scans revealed hypometabolic areas of the olfactory gyrus, amygdala, hippocampus, cingulum and pons (11). We also recently showed that in a patient who had a mild COVID infection (without hospitalization or oxygen supplementation), a marked attentional and executive cognitive impairment was associated on FDG PET with hypometabolic areas of the cingulate cortex (12). This imaging exam represents an important diagnostic tool for visualizing post-COVID brain damage in subjects with cognitive complaints (brain fog) and confirming that long COVID can be linked to neurological sequelae.

The long-term course of these brain lesions and clinical symptoms in mild forms of COVID is difficult to predict. Some authors have mentioned that an evolution towards neurodegenerative diseases could be seen over a prolonged evolution (2). It has been reported in previous studies that SARS-CoV-2 can induce hyperphosphorylation of tau protein or activation of the PKR kinase, two proteins that abnormally accumulate in the brains of Alzheimer's patients (12, 13). Long-term neurological and cognitive monitoring of these patients seems to be an important measure in terms of public health and a new approach toward neurodegenerative diseases.

It seems useful to introduce sustained cognitive remediation for these patients; in the event of an associated depression, serotonin reuptake inhibitors, which have an anti-inflammatory effect in the brain, could be proposed (14)

In conclusion, the cognitive disorders associated with long COVID following mild forms of infection could be linked to hypometabolic lesions of the brain, which are probably associated with neuroinflammation. Only long-term follow-up of the patients will be able to conclusively assess the evolution of these lesions and the accompanying clinical signs.

To sum up, neurological complications of COVID may be detected in moderately affected individuals and extensive explorations of these patients should be performed to determine the actual extension of the brain lesions.

1. Wiersinga WJ, Rhodes A, Cheng AC, Peacock SJ, Prescott HC. Pathophysiology, Transmission, Diagnosis, and Treatment of Coronavirus Disease 2019 (COVID-19): A Review. *JAMA*. 2020;324(8):782-93.
2. Heneka MT, Golenbock D, Latz E, Morgan D, Brown R. Immediate and long-term consequences of COVID-19 infections for the development of neurological disease. *Alzheimers Res Ther*. 2020;12(1):69.
3. Thakur KT, Miller EH, Glendinning MD, Al-Dalahmah O, Banu MA, Boehme AK, et al. COVID-19 neuropathology at Columbia University Irving Medical Center/New York Presbyterian Hospital. *Brain*. 2021.
4. Serrano GE, Walker JE, Arce R, Glass MJ, Vargas D, Sue LI, et al. Mapping of SARS-CoV-2 Brain Invasion and Histopathology in COVID-19 Disease. *medRxiv*. 2021.
5. Virhammar J, Naas A, Fallmar D, Cunningham JL, Klang A, Ashton NJ, et al. Biomarkers for central nervous system injury in cerebrospinal fluid are elevated in COVID-19 and associated with neurological symptoms and disease severity. *Eur J Neurol*. 2020.
6. Taquet M, Geddes JR, Husain M, Luciano S, Harrison PJ. 6-month neurological and psychiatric outcomes in 236 379 survivors of COVID-19: a retrospective cohort study using electronic health records. *Lancet Psychiatry*. 2021;8(5):416-27.
7. Helms J, Kremer S, Merdji H, Clere-Jehl R, Schenck M, Kummerlen C, et al. Neurologic Features in Severe SARS-CoV-2 Infection. *N Engl J Med*. 2020;382(23):2268-70.
8. Ellul MA, Benjamin L, Singh B, Lant S, Michael BD, Easton A, et al. Neurological associations of COVID-19. *Lancet Neurol*. 2020;19(9):767-83.
9. Nalbandian A, Sehgal K, Gupta A, Madhavan MV, McGroder C, Stevens JS, et al. Post-acute COVID-19 syndrome. *Nat Med*. 2021;27(4):601-15.

10. Graham EL, Clark JR, Orban ZS, Lim PH, Szymanski AL, Taylor C, et al. Persistent neurologic symptoms and cognitive dysfunction in non-hospitalized Covid-19 "long haulers". *Ann Clin Transl Neurol.* 2021;8(5):1073-85.
11. Guedj E, Million M, Dudouet P, Tissot-Dupont H, Bregeon F, Cammilleri S, et al. (18)F-FDG brain PET hypometabolism in post-SARS-CoV-2 infection: substrate for persistent/delayed disorders? *Eur J Nucl Med Mol Imaging.* 2021;48(2):592-5.
12. Hugon J, Msika EF, Queneau M, Farid K, Paquet C. Long COVID: cognitive complaints (brain fog) and dysfunction of the cingulate cortex. *J Neurol.* 2021.
13. Ramani A, Muller L, Ostermann PN, Gabriel E, Abida-Islam P, Muller-Schiffmann A, et al. SARS-CoV-2 targets neurons of 3D human brain organoids. *EMBO J.* 2020;39(20):e106230.
14. Zhang X, Zhu LB, He JH, Zhang HQ, Ji SY, Zhang CN, et al. Paroxetine suppresses reactive microglia-mediated but not lipopolysaccharide-induced inflammatory responses in primary astrocytes. *J Neuroinflammation.* 2020;17(1):50.